

Taking Joenja, the **first** and **only FDA-approved** treatment for APDS, is as easy as...

1 70 mg tablet **2** times a day **3** ways to make sure every dose counts

Establish a treatment routine with Joenja



Take your Joenja tablets about **12 hours apart** at typically convenient times



Set **2 reminders**, one in the morning and another at night



Keep your Joenja tablets **in the same place** so you always know where they are

Joenja
leniolisib
HOPE UPLIFTED

Approved Use

JOENJA® (leniolisib) is a prescription medicine that is used to treat activated phosphoinositide 3-kinase delta (PI3Kδ) syndrome (APDS) in adults and children 12 years of age and older.

Select Safety Information

Tell your healthcare provider if you are pregnant or plan to become pregnant. JOENJA may harm your unborn baby. Your healthcare provider will do a pregnancy test before you start receiving JOENJA.

Use effective birth control to prevent pregnancy while taking JOENJA and for one week after you stop taking JOENJA. If you plan to become pregnant, continue taking your birth control for one week after stopping JOENJA. Talk to your doctor about what type of birth control method is right for you while taking JOENJA.

Please see additional Select Safety Information throughout.

For more information, please see the full [Prescribing Information](#).

Getting started with Joenja

Joenja
leniolisib
HOPE UPLIFTED

How do I take Joenja?

Joenja is an oral tablet, taken twice a day, anytime, anywhere.

- Take Joenja exactly as your doctor tells you to take it
- Take 1 Joenja tablet in the morning, wait about 12 hours, and take the second tablet at night. Joenja can be taken with or without food
- If you miss a dose of Joenja by more than 6 hours, take the next dose at your usual time
- If vomiting occurs **within 1 hour** after taking Joenja, take another tablet as soon as you are able
- If vomiting occurs **more than 1 hour** after dosing, wait and take the next dose at the usual time

Never miss a dose!

Download the free Joenja reminder tone today at [Joenja.com](https://www.joenja.com)

Select Safety Information (cont.)

It is not known if JOENJA passes into your breast milk. Talk to your doctor about the best way to feed your baby if you take JOENJA.

Please see additional Select Safety Information throughout.

For more information, please see the full [Prescribing Information](#).



Not an actual patient.

Do I need to keep taking Joenja?

- **DO NOT** stop taking Joenja without talking to your doctor. Always speak with your doctor before making any treatment changes or decisions
- Take Joenja twice a day, every day, to keep your APDS under control even if it makes you feel better

Can I take Joenja with my other medications?

- Joenja and certain other medicines can interact with each other. Tell your doctor about the medicines you take. They may affect how Joenja works

This information is intended to help you have the best possible experience while taking Joenja. This guide does not take the place of talking to your doctor about your medical condition or your treatment.

Select Safety Information (cont.)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. JOENJA and certain other medicines may affect each other.

Tell your healthcare provider if you are scheduled to receive an immunization (vaccine). JOENJA may affect how well a vaccine works.

Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.

Please see additional Select Safety Information throughout.

For more information, please see the full Prescribing Information.



Select Safety Information (cont.)

The most common side effects of **JOENJA** include headache, inflammation of sinuses, and dry, itchy, and inflamed skin (eczema). These are not all the possible side effects of JOENJA. Tell your healthcare provider about any side effect that bothers you or does not go away.

You can also report negative side effects to FDA at 1-800-FDA-1088 or www.fda.gov/medwatch, or Pharming Medical Affairs at (800) 930-5221.

Please see additional Select Safety Information throughout.

For more information, please see the full [Prescribing Information](#).

Not actual patients.



What if I need help getting Joenja?

APDS Assist is here to help you navigate your treatment needs.

To connect with a Care Coordinator at APDS Assist, call 1-877-796-APDS (2737) between 8 AM-8 PM ET, Monday-Friday.

Visit Joenja.com for more information and ask your doctor about enrollment.



Focus on the
things you love.

Joenja can help get your
APDS under control.



Joenja and logo are registered trademarks of Pharming Intellectual Property, B.V.
©2023 Pharming Healthcare, Inc. All rights reserved.
JOE-US-2023-0074 03/23